

Abelian varieties associated to Gaussian lattices

Arnaud Beauville

To Joe Harris on his 60th birthday

ABSTRACT. We associate to a unimodular lattice Γ , endowed with an automorphism i of square -1 , a principally polarized abelian variety $A_\Gamma = \Gamma_{\mathbb{R}}/\Gamma$. We show that the configuration of i -invariant theta divisors of A_Γ follows a pattern very similar to the classical theory of theta characteristics; as a consequence we find that A_Γ has a high number of vanishing thetanulls. When $\Gamma = E_8$ we recover the 10 vanishing thetanulls of the abelian fourfold discovered by R. Varley.

Introduction

A *Gaussian lattice* is a free, finitely generated $\mathbb{Z}[i]$ -module Γ with a positive hermitian form $\Gamma \times \Gamma \rightarrow \mathbb{Z}[i]$. Equivalently, we can view Γ as a lattice over \mathbb{Z} endowed with an automorphism i of square -1_Γ . This gives a complex structure on the vector space $\Gamma_{\mathbb{R}} := \Gamma \otimes_{\mathbb{Z}} \mathbb{R}$; we associate to Γ the complex torus $A_\Gamma := \Gamma_{\mathbb{R}}/\Gamma$.

As a complex torus A_Γ is isomorphic to E^g , where E is the complex elliptic curve $\mathbb{C}/\mathbb{Z}[i]$ and $g = \frac{1}{2} \text{rk}_{\mathbb{Z}} \Gamma$. More interestingly, the hermitian form provides a *polarization* on A_Γ (see (1.3) below); in particular, if Γ is unimodular, A is a principally polarized abelian variety (p.p.a.v. for short), which is indecomposable if Γ is indecomposable.

The first non-trivial case is $g = 4$, with Γ the root lattice of type E_8 (Example 1.2.1). The resulting p.p.a.v. is the abelian fourfold discovered by Varley [V] with a different (and more geometric) description; it has 10 “vanishing thetanulls” (even theta functions vanishing at 0), the maximum possible for a 4-dimensional indecomposable p.p.a.v. In fact this property characterizes the Varley fourfold outside the hyperelliptic Jacobian locus [D].

Our aim is to explain this property from the lattice point of view, and to extend it to all unimodular lattices. It turns out that we can mimic the classical theory of theta characteristics, replacing the automorphism (-1) by i . We will show:

- The group A_i of i -invariant points of A_Γ is a vector space of dimension g over $\mathbb{Z}/2$; it admits a natural non-degenerate bilinear symmetric form b .

- The set of i -invariant theta divisors of A_Γ is an affine space over A_i , isomorphic to the space of quadratic forms on A_i associated to b (see (2.1)).
- Let Θ be an i -invariant theta divisor, and Q the corresponding quadratic form. The multiplicity $m_0(\Theta)$ of Θ at 0 satisfies

$$2m_0(\Theta) \equiv \sigma(Q) + g \pmod{8},$$

where σ is the *Brown invariant* of the form Q (2.1).

As a consequence, we obtain a high number of i -invariant divisors Θ with $m_0(\Theta) \equiv 2 \pmod{4}$; each of them corresponds to a vanishing thetanull. When Γ is even, this number is $2^{\frac{g}{2}-1}(2^{\frac{g}{2}} - (-1)^{\frac{g}{4}})$; for $g = 4$ we recover the 10 vanishing thetanulls of the Varley fourfold.

1. Gaussian lattices

1.1. Lattices. As recalled in the Introduction, a Gaussian lattice is a free finitely generated $\mathbb{Z}[i]$ -module Γ endowed with a positive hermitian form¹ $H : \Gamma \times \Gamma \rightarrow \mathbb{Z}[i]$. We write $H(x, y) = S(x, y) + iE(x, y)$; S and E are \mathbb{Z} -bilinear forms on Γ , S is symmetric, E is skew-symmetric, and we have

$$S(ix, iy) = S(x, y) \quad , \quad E(ix, iy) = E(x, y) \quad , \quad E(x, y) = S(ix, y) .$$

We will rather view a Gaussian lattice as an ordinary lattice (over \mathbb{Z}) with an automorphism i such that $i^2 = -1_\Gamma$: the last formula above defines E , and we have $H = S + iE$.

We have $\det S = \det E = (\det H)^2$; the lattice is *unimodular* when these numbers are equal to 1. It is *even* if $S(x, x)$ is even for all $x \in \Gamma$. We say that Γ is *indecomposable over $\mathbb{Z}[i]$* if it cannot be written as the orthogonal sum of two nonzero Gaussian lattices; this is of course the case if Γ is indecomposable over \mathbb{Z} , but the converse is false (Example 3 below).

1.2. Examples. 1) For g even, the lattice Γ_{2g} is

$$\Gamma_{2g} := \{(x_j) \in \mathbb{R}^{2g} \mid x_j \in \frac{1}{2}\mathbb{Z}, x_j - x_k \in \mathbb{Z}, \sum x_j \in 2\mathbb{Z}\} .$$

The inner product is inherited from the euclidean structure of \mathbb{R}^{2g} , and the automorphism i is given in the standard basis (e_j) by

$$ie_{2j-1} = e_{2j} \quad ie_{2j} = -e_{2j-1} \quad \text{for } 1 \leq j \leq g .$$

The lattice Γ_{2g} is unimodular, indecomposable when $g > 2$, and even if g is divisible by 4. The first case $g = 4$ gives the root lattice E_8 .

The automorphism i is *unique* up to conjugacy: for $g = 4$ this is classical [C], and for $g \geq 6$ this follows easily from the fact that $\text{Aut}(\Gamma_{2g})$ is the semi-direct product $(\mathbb{Z}/2)^{2g-1} \rtimes \mathfrak{S}_{2g}$, acting by permutation and even changes of sign of the basis vectors (e_j) .

¹Our convention is that $H(x, y)$ is \mathbb{C} -linear in y .

2) The Leech lattice Λ_{24} admits an automorphism of square -1 [**C-S**], also unique up to conjugacy.

3) Let Γ_0 be a lattice, and $\Gamma := \Gamma_0 \otimes_{\mathbb{Z}} \mathbb{Z}[i]$. The inner product of Γ_0 extends to an hermitian inner product on Γ , which is then a gaussian lattice. If Γ_0 is unimodular, resp. even, resp. indecomposable, Γ is unimodular, resp. even, resp. indecomposable over $\mathbb{Z}[i]$.

1.3. The abelian variety A_{Γ} . Let Γ be a Gaussian lattice, of rank $2g$ over \mathbb{Z} . We put $\Gamma_{\mathbb{R}} := \Gamma \otimes_{\mathbb{Z}} \mathbb{R}$ and $A_{\Gamma} := \Gamma_{\mathbb{R}}/\Gamma$. The automorphism i defines a complex structure on $\Gamma_{\mathbb{R}}$, so that A_{Γ} is a complex torus. Since Γ is a free $\mathbb{Z}[i]$ -module, A_{Γ} is isomorphic to E^g , where E is the complex elliptic curve $\mathbb{C}/\mathbb{Z}[i]$.

The positive hermitian form H extends to $\Gamma_{\mathbb{R}}$, and its imaginary part E takes integral values on Γ : this is by definition a *polarization* on A_{Γ} . The polarization is principal if and only if Γ is unimodular; the p.p.a.v. A_{Γ} is indecomposable (i.e. is not a product of two nontrivial p.p.a.v.) if and only if Γ is indecomposable over $\mathbb{Z}[i]$.

The multiplication by i on $\Gamma_{\mathbb{R}}$ induces an automorphism of A_{Γ} , that we simply denote i . Conversely, let $A = V/\Gamma$ be a complex torus, of dimension g , with an automorphism inducing on $T_0(A) = V$ the multiplication by i . Then Γ is a $\mathbb{Z}[i]$ -module, thus isomorphic to $\mathbb{Z}[i]^g$, so that A is isomorphic to E^g ; polarizations of A correspond bijectively to positive hermitian forms on Γ .

2. Linear algebra over $(\mathbb{Z}/2)[i]$

2.1. Linear algebra over $\mathbb{Z}/2$. We consider a vector space V over $\mathbb{Z}/2$, of dimension g , with a non-degenerate symmetric bilinear form b on V . The form $x \mapsto b(x, x)$ is linear. Two different situations may occur:

- $b(x, x) = 0$ for all $x \in V$; in that case b is a symplectic form.
- $b(x, x)$ is not identically zero; it is then easy (using induction on g) to prove that V admits an orthonormal basis with respect to b .

A *quadratic form associated to b* is a function $q : V \rightarrow \mathbb{Z}/4$ such that

$$q(x + y) = q(x) + q(y) + 2b(x, y) \quad \text{for } x, y \in V,$$

where multiplication by 2 stands for the isomorphism $\mathbb{Z}/2 \xrightarrow{\sim} 2\mathbb{Z}/4\mathbb{Z} \subset \mathbb{Z}/4\mathbb{Z}$.

Observe that this implies $q(0) = 0$ and $q(x) \equiv b(x, x) \pmod{2}$. We denote by \mathcal{Q}_b the set of quadratic forms associated to b ; \mathcal{Q}_b is an affine space over V , the action of V being given by $(\alpha + q)(x) = q(x) + 2b(\alpha, x)$ for $q \in \mathcal{Q}_b$, $\alpha, x \in V$.

When b is symplectic, q takes its values in $2\mathbb{Z}/4\mathbb{Z} \cong \mathbb{Z}/2$; the corresponding form $q' : V \rightarrow \mathbb{Z}/2$ is a quadratic form associated to b in the usual sense, that is satisfies $q'(x + y) = q'(x) + q'(y) + b(x, y)$ for $x, y \in V$.

The *Brown invariant* $\sigma(q) \in \mathbb{Z}/8$ of a form $q \in \mathcal{Q}_b$ has been introduced in [**B**] as a generalization of the Arf invariant; it can be defined as follows. If b is symplectic, we put $\sigma(q) := 4\text{Arf}(q')$, where $q' : V \rightarrow \mathbb{Z}/2$ is the form defined

above. Otherwise b admits an orthonormal basis (e_1, \dots, e_g) ; we have $q(e_i) = \pm 1$, and we let g^+ (resp. g^-) be the number of basis vectors e_i such that $q(e_i) = 1$ (resp. -1). Then $\sigma(q) = g^+ - g^- \pmod{8}$.

2.2. Linear algebra over $(\mathbb{Z}/2)[i]$. Let Γ be a unimodular Gaussian lattice of rank $2g$ over \mathbb{Z} . We put $A_2 := \Gamma/2\Gamma$; this is naturally identified with the 2-torsion subgroup of A_Γ . We have the following structures on A_2 :

a) A_2 is a free $(\mathbb{Z}/2)[i]$ -module of rank g . We put $\varepsilon := 1 + i$ in $(\mathbb{Z}/2)[i]$; then $(\mathbb{Z}/2)[i] = (\mathbb{Z}/2)[\varepsilon]$, with $\varepsilon^2 = 0$. The subgroup A_i of i -invariant elements is $\text{Ker } \varepsilon = \varepsilon A_2$; it is a vector space of dimension g over $\mathbb{Z}/2$.

b) The form E induces on A_2 a symplectic form e (the Weil pairing for A_Γ). Since $E(x, iy) = -E(ix, y)$, we have, for $\alpha, \beta \in A_2$,

$$e(\alpha, \varepsilon\beta) = e(\varepsilon\alpha, \beta) \quad \text{hence} \quad e(\varepsilon\alpha, \varepsilon\beta) = 0;$$

thus A_i is a Lagrangian subspace of A_2 .

c) The form $x \mapsto S(x, x)$ induces a quadratic form $Q : A_2 \rightarrow \mathbb{Z}/4$ associated with the bilinear symmetric form $(\alpha, \beta) \mapsto e(\alpha, i\beta)$ (2.1). In particular we have $Q(\alpha) \equiv e(\alpha, i\alpha) \pmod{2}$.

Since $S((1+i)x, (1+i)x) = 2S(x, x)$, we have $Q(\varepsilon\alpha) = 2Q(\alpha) = 2e(\alpha, i\alpha)$.

LEMMA 1. *Let $q : A_2 \rightarrow \mathbb{Z}/4$ be an i -invariant quadratic form associated to e . The formulas*

$$b(\varepsilon\alpha, \varepsilon\beta) = e(\alpha, \varepsilon\beta) \quad , \quad Q_q(\varepsilon\alpha) = q(\alpha) - Q(\alpha) \quad \text{for } \alpha, \beta \in A_2,$$

define on $A_i = \varepsilon A_2$ a non-degenerate symmetric form b and a quadratic form $Q_q : A_i \rightarrow \mathbb{Z}/4$ associated with b .

Proof : Since $A_i = \text{Ker } \varepsilon$ is isotropic for e , the expression $e(\alpha, \varepsilon\beta)$ is a bilinear function b of $\varepsilon\alpha$ and $\varepsilon\beta$; it is symmetric by b). If $e(\alpha, \varepsilon\beta) = 0$ for all β in A_2 we have $\alpha \in A_i$ because A_i is Lagrangian, hence $\varepsilon\alpha = 0$, so b is non-degenerate.

Put $\tilde{Q}_q(\alpha) = q(\alpha) - Q(\alpha) \in \mathbb{Z}/4$ for $\alpha \in A_2$. We have

$$\tilde{Q}_q(\alpha + \beta) = \tilde{Q}_q(\alpha) + \tilde{Q}_q(\beta) + 2e(\alpha, \varepsilon\beta).$$

Take $\beta = \varepsilon\gamma$. Since q is i -invariant we have $q(\varepsilon\gamma) = 2e(\gamma, i\gamma) = Q(\varepsilon\gamma)$ by c), hence $\tilde{Q}_q(\varepsilon\gamma) = 0$ and $\tilde{Q}_q(\alpha + \varepsilon\gamma) = \tilde{Q}_q(\alpha)$. Thus \tilde{Q}_q defines a quadratic form Q_q on A_i associated to b . \blacksquare

Let $\mathcal{Q}_e^{(i)}$ be the set of i -invariants quadratic forms on A_2 associated to e . If $q \in \mathcal{Q}_e^{(i)}$ and $\alpha \in A_2$, we have $\alpha + q \in \mathcal{Q}_e^{(i)}$ if and only if α belongs to $A_i^\perp = A_i$; in other words, $\mathcal{Q}_e^{(i)}$ is an affine subspace of \mathcal{Q}_e , with direction A_i .

LEMMA 2. *The map $q \mapsto Q_q$ is an affine isomorphism of $\mathcal{Q}_e^{(i)}$ onto \mathcal{Q}_b .*

Proof : We just have to prove the equality $Q_{\alpha+q} = \alpha + Q_q$ for $q \in \mathcal{Q}_e^{(i)}$, $\alpha \in A_i$. Let $\beta \in A_i$; we write $\beta = \varepsilon\beta'$ for some $\beta' \in A_2$. Then

$$Q_{\alpha+q}(\beta) = 2e(\alpha, \beta') + q(\beta') - Q(\beta') = 2b(\alpha, \beta) + Q_q(\beta). \quad \blacksquare$$

REMARK 1. Let $\alpha \in A_2$; we have $b(\varepsilon\alpha, \varepsilon\alpha) = e(\alpha, \varepsilon\alpha) = e(\alpha, i\alpha) \equiv Q(\alpha) \pmod{2}$, hence *the form b is symplectic if and only if Γ is even*. In this case we have $e(\alpha, i\alpha) = 0$ for all $\alpha \in A_2$; it follows that $\mathcal{Q}_e^{(i)}$ is the set of forms vanishing on A_i . Since A_i is Lagrangian for e , this implies that these forms, viewed as quadratic forms $A_2 \rightarrow \mathbb{Z}/2$, are all even (that is, their Arf invariant is 0).

3. i -invariant theta divisors

3.1. Reminder on theta characteristics. We first recall the classical theory of theta characteristics on an arbitrary p.p.a.v. $A = V/\Gamma$. Let $A_2 \cong \Gamma/2\Gamma$ be the 2-torsion subgroup of A , \mathcal{T} the set of symmetric theta divisors on A , and \mathcal{Q}_e the set of quadratic forms on A_2 associated to the Weyl pairing e . The $\mathbb{Z}/2$ -vector space A_2 acts on \mathcal{T} by translation, and on \mathcal{Q}_e by the action defined in (2.1); both sets are affine spaces over A_2 , and there is a canonical affine isomorphism $q \mapsto \Theta_q$ of \mathcal{Q}_e onto \mathcal{T} . It can be defined as follows ([M], §2). Let $\gamma \in \Gamma$, and let $\bar{\gamma}$ be its class in A_2 . For $z \in V$, we put

$$e_\gamma(z) = i^{q(\bar{\gamma})} e^{\pi H(\gamma, z + \frac{\gamma}{2})}.$$

We define an action of Γ on the trivial bundle $V \times \mathbb{C}$ by $\gamma.(z, t) = (z + \gamma, e_\gamma(z)t)$; then the quotient of $V \times \mathbb{C}$ by this action is the line bundle $\mathcal{O}_A(\Theta_q)$ on A .

3.2. The main results. We go back to the abelian variety A_Γ associated to a Gaussian lattice Γ . We assume that Γ is unimodular. We use the notation of (2.2). The isomorphism $\mathcal{Q}_e \xrightarrow{\sim} \mathcal{T}$ is compatible with the action of i , so i -invariant theta divisors correspond to forms $q \in \mathcal{Q}_e^{(i)}$.

Let $q \in \mathcal{Q}_e^{(i)}$, and let L be the line bundle $\mathcal{O}_{A_\Gamma}(\Theta_q)$. We have $i^*L \cong L$; we denote by $\iota : i^*L \rightarrow L$ the unique isomorphism inducing the identity of L_0 . For each $\alpha \in A_i$, ι induces an isomorphism $\iota(\alpha) : L_\alpha \rightarrow L_\alpha$.

PROPOSITION 1. $\iota(\alpha)$ is the homothety of ratio $i^{Q_q(\alpha)}$.

Proof : The isomorphism $\iota^{-1} : L \xrightarrow{\sim} i^*L$ corresponds to a linear automorphism j of L above i :

$$\begin{array}{ccc} L & \xrightarrow{j} & L \\ \downarrow & & \downarrow \\ A_\Gamma & \xrightarrow{i} & A_\Gamma \end{array}.$$

Consider the automorphism $\tilde{j} : (z, t) \mapsto (iz, t)$ of $\Gamma_{\mathbb{R}} \times \mathbb{C}$. Since $e_{i\gamma}(iz) = e_\gamma(z)$, we have $\tilde{j}(\gamma.(z, t)) = (i\gamma).\tilde{j}(z, t)$. Thus \tilde{j} factors through an isomorphism $L \rightarrow L$

above i which is the identity on L_0 , hence equal to j ; that is, we have a commutative diagram:

$$\begin{array}{ccc} \Gamma_{\mathbb{R}} \times \mathbb{C} & \xrightarrow{\tilde{j}} & \Gamma_{\mathbb{R}} \times \mathbb{C} \\ \pi \downarrow & & \downarrow \pi \\ L & \xrightarrow{j} & L \end{array}$$

where π is the quotient map.

Let $\alpha \in A_i$, and let γ be an element of Γ whose class (mod. 2Γ) is α . Then $\delta := \frac{i\gamma}{2} - \frac{\gamma}{2}$ belongs to Γ . We have

$$j(\pi(\frac{\gamma}{2}, t)) = \pi(\frac{i\gamma}{2}, t) = \pi(\frac{\gamma}{2}, e_{\delta}(\frac{\gamma}{2})^{-1}t),$$

hence $\iota(\alpha) = j(\alpha)^{-1}$ is the homothety of ratio $e_{\delta}(\frac{\gamma}{2})$. Let β be the class of δ in A_2 . Since $\gamma = -(1+i)\delta$, we have $\alpha = \varepsilon\beta$, hence

$$e_{\delta}(\frac{\gamma}{2}) = i^{q(\beta)} e^{\frac{\pi}{2}H(\delta, \gamma + \delta)} = i^{q(\beta) - H(\delta, \delta)} = i^{Q_q(\alpha)}. \quad \blacksquare$$

From $\iota : i^*L \rightarrow L$ we deduce an isomorphism $\iota^b : L \xrightarrow{\sim} i_*L$, inducing on global sections an automorphism of $H^0(A_{\Gamma}, L)$.

PROPOSITION 2. ι^b acts on $H^0(A_{\Gamma}, L)$ by multiplication by $e^{\frac{i\pi}{4}(\sigma(Q_q) + g)}$.

Note that $\sigma(Q_q) \equiv g \pmod{2}$ ([B], Thm. 1.20, (vi)), so this number is a power of i .

Proof : Since $\dim H^0(A_{\Gamma}, L) = 1$ it suffices to compute $\text{Tr } \iota^b$. This is given by the holomorphic Lefschetz formula [A-B] applied to (i, ι) . Since $H^i(A_{\Gamma}, L) = 0$ for $i > 0$, we find

$$\text{Tr } \iota^b = \sum_{\alpha \in A_i} \frac{\text{Tr } \iota(\alpha)}{(1-i)^g} = (1-i)^{-g} \sum_{\alpha \in A_i} i^{Q_q(\alpha)}.$$

We have $(1-i)^{-g} = 2^{-\frac{g}{2}} e^{\frac{i\pi g}{4}}$ and $\sum_{\alpha \in A_i} i^{Q_q(\alpha)} = 2^{\frac{g}{2}} e^{\frac{i\pi}{4}\sigma(Q_q)}$ ([B], Thm. 1.20, (xi)), hence the result. \blacksquare

PROPOSITION 3. Let $\alpha \in A_i$, and let $m_{\alpha}(\Theta_q)$ be the multiplicity of Θ_q at α . We have

$$2m_{\alpha}(\Theta_q) \equiv \sigma(Q_q) + g - 2Q_q(\alpha) \pmod{8}.$$

Proof : Let θ be a nonzero section of $H^0(A_{\Gamma}, L)$. Choose a local non-vanishing section s of L around α . We can write $\theta = fs$ in a neighborhood of α , with $f \in \mathcal{O}_{A_{\Gamma}, \alpha}$. We have $\iota^b(\theta) = i^k \theta$ with $2k \equiv \sigma(Q_q) + g \pmod{8}$ (Proposition 2), hence

$$(i^*f)\iota^b(s) = i^k fs.$$

We look at this equality in $\mathfrak{m}_{\alpha}^m L / \mathfrak{m}_{\alpha}^{m+1} L$, where \mathfrak{m}_{α} is the maximal ideal of $\mathcal{O}_{A_{\Gamma}, \alpha}$ and $m := m_{\alpha}(\Theta)$. We have $i^*f = i^m f \pmod{\mathfrak{m}_{\alpha}^{m+1}}$, and $\iota^b(s) = \iota(\alpha)s \pmod{\mathfrak{m}_{\alpha} L}$. We obtain $i^m \iota(\alpha) = i^k$, hence the result in view of Proposition 1. \blacksquare

COROLLARY. *The number of i -invariant theta divisors Θ with $m_0(\Theta) \equiv 2 \pmod{4}$ is*

$$2^{\frac{g}{2}-1}(2^{\frac{g}{2}} - (-1)^{\frac{g}{4}}) \quad \text{if } \Gamma \text{ is even, and } \quad 2^{g-2} - 2^{\frac{g}{2}-1} \cos \frac{\pi g}{4} \quad \text{if } \Gamma \text{ is odd;}$$

each of these divisors corresponds to a vanishing thetanull.

Proof : According to the Proposition, we have $m_0(\Theta_q) \equiv 2 \pmod{4}$ if and only if $\sigma(Q_q) \equiv 4 - g \pmod{8}$. When q runs over $\mathcal{Q}_e^{(i)}$, Q_q runs over \mathcal{Q}_b (Lemma 2.2), so we must find how many elements Q of \mathcal{Q}_b satisfy $\sigma(Q) \equiv 4 - g \pmod{8}$.

If Γ is even (so that g is divisible by 4), we identify \mathcal{Q}_b with the set of quadratic forms $Q : A_2 \rightarrow \mathbb{Z}/2$ associated with the symplectic form b ; the previous congruence becomes $\text{Arf}(Q) \equiv 1 + \frac{g}{4} \pmod{2}$. There are $2^{\frac{g}{2}-1}(2^{\frac{g}{2}} + 1)$ such forms with Arf invariant 0 and $2^{\frac{g}{2}-1}(2^{\frac{g}{2}} - 1)$ with Arf invariant 1, hence the result.

Assume that Γ is odd; we choose an orthonormal basis (e_1, \dots, e_g) for b . The forms $Q \in \mathcal{Q}_b$ are determined by their values $Q(e_i) = \pm 1$; the condition is that the number g^+ of +1 values satisfies

$$2g^+ - g \equiv 4 - g \pmod{8}, \quad \text{hence } g^+ \equiv 2 \pmod{4}.$$

The number of forms with the required property is thus the number of subsets $E \subset \{1, \dots, g\}$ with $\text{Card}(E) \equiv 2 \pmod{4}$, that is

$$\binom{g}{2} + \binom{g}{6} + \dots = \frac{1}{4} [(1+1)^g + (1-1)^g - (1+i)^g - (1-i)^g] = 2^{g-2} - 2^{\frac{g}{2}-1} \cos \frac{\pi g}{4}. \quad \blacksquare$$

Thus we find a number of vanishing thetanulls asymptotically equivalent to 2^{g-1} when Γ is even, and 2^{g-2} when Γ is odd. These numbers are rather modest, at least by comparison with the number of vanishing thetanulls of a hyperelliptic Jacobian, which is asymptotically equivalent to 2^{2g-1} . However, when Γ is even, the vanishing thetanulls of A_Γ have the particular property of being ‘‘syzygetic’’ in the classical terminology, which just means that the corresponding quadratic forms (3.1) lie in an affine subspace of \mathcal{Q}_e which consists of even forms (Remark 1). Such a subspace has dimension $\leq g$, and it might be that the number given by the Corollary in the even case is the maximum possible for a syzygetic subset of vanishing thetanulls.

4. Complements

4.1. Automorphisms. The automorphism group of A_Γ is the centralizer of i in $\text{Aut}(\Gamma)$. This group can be rather large: it has order 46080 for $\Gamma = E_8$ and 2012774400 for $\Gamma = \Lambda_{24}$ [C-S]. For the lattice Γ_{2g} (Example 1.2.1) with $g > 4$, it has order $2^{2g-1}g!$.

For the lattice $\Gamma = \Gamma_0 \otimes_{\mathbb{Z}} \mathbb{Z}[i]$ of Example 1.2.3, $\text{Aut}(A_\Gamma)$ is generated by i and the group $\text{Aut}(\Gamma_0)$. Note that there are examples of unimodular lattices (even or odd) Γ_0 with $\text{Aut}(\Gamma_0) = \{\pm 1\}$ [Ba], so that $\text{Aut}(A_\Gamma)$ is reduced to $\{\pm 1, \pm i\}$.

4.2. Jacobians. We observe that for $g > 1$ the p.p.a.v. A_Γ can *not* be a Jacobian. Indeed, let C be a curve of genus g ; if $JC \cong A_\Gamma$, Torelli theorem provides an automorphism u of C inducing either i or $-i$ on JC , hence also on $T_0(JC) = H^0(C, K_C)^*$. Then u acts trivially on the image of the canonical map $C \rightarrow \mathbb{P}(H^0(C, K_C)^*)$; this implies that u is the identity or that C is hyperelliptic and u is the hyperelliptic involution. But in these cases u acts on $H^0(C, K_C)$ by multiplication by ± 1 , a contradiction. ■

References

- [A-B] M. Atiyah, R. Bott : *A Lefschetz Fixed Point Formula for Elliptic Complexes: II. Applications*. Ann. of Math. (2) **88** (1968), 451–491. MR0232406 (38:731)
- [Ba] R. Bacher : *Unimodular lattices without nontrivial automorphisms*. Int. Math. Res. Notes **2** (1994), 91–95. MR1264932 (95b:11067)
- [B] E.H. Brown, Jr. : *Generalizations of the Kervaire invariant*. Ann. of Math. (2) **95** (1972), 368–383. MR0293642 (45:2719)
- [C] R. W. Carter : *Conjugacy classes in the Weyl group*. Compositio Math. **25** (1972), 1–59. MR0318337 (47:6884)
- [C-S] J.H. Conway, N. Sloane : *D_4 , E_8 , Leech and certain other lattices are symplectic*. Invent. Math. **117** (1994), no. 1, 53–55.
- [D] O. Debarre : *Annulation de thétaconstantes sur les variétés abéliennes de dimension quatre*. C. R. Acad. Sci. Paris Sér. I Math. **305** (1987), no. 20, 885–888. MR925290 (89a:14055)
- [M] D. Mumford: *Abelian varieties*. Oxford University Press, London, 1970. MR0282985 (44:219)
- [V] R. Varley : *Weddle’s surfaces, Humbert’s curves, and a certain 4-dimensional abelian variety*. Amer. J. Math. **108** (1986), no. 4, 931–952. MR853219 (87g:14050)

LABORATOIRE J.-A. DIEUDONNÉ, UMR 7351 DU CNRS, UNIVERSITÉ DE NICE, PARC VALROSE,
F-06108 NICE CEDEX 2, FRANCE

E-mail address: arnaud.beauville@unice.fr