

# SEMISTABILITY OF FROBENIUS DIRECT IMAGES OVER CURVES

VIKRAM B. MEHTA AND CHRISTIAN PAULY

ABSTRACT. Let  $X$  be a smooth projective curve of genus  $g \geq 2$  defined over an algebraically closed field  $k$  of characteristic  $p > 0$ . Given a semistable vector bundle  $E$  over  $X$ , we show that its direct image  $F_*E$  under the Frobenius map  $F$  of  $X$  is again semistable. We deduce a numerical characterization of the stable rank- $p$  vector bundles  $F_*L$ , where  $L$  is a line bundle over  $X$ .

## 1. INTRODUCTION

Let  $X$  be a smooth projective curve of genus  $g \geq 2$  defined over an algebraically closed field  $k$  of characteristic  $p > 0$  and let  $F : X \rightarrow X_1$  be the relative  $k$ -linear Frobenius map. It is by now a well-established fact that on any curve  $X$  there exist semistable vector bundles  $E$  such that their pull-back under the Frobenius map  $F^*E$  is not semistable [LanP], [LasP]. In order to control the degree of instability of the bundle  $F^*E$ , one is naturally lead (through adjunction  $\mathrm{Hom}_{\mathcal{O}_X}(F^*E, E') = \mathrm{Hom}_{\mathcal{O}_{X_1}}(E, F_*E')$ ) to ask whether semistability is preserved by direct image under the Frobenius map. The answer is (somewhat surprisingly) yes. In this note we show the following result.

**1.1. Theorem.** *Assume that  $g \geq 2$ . If  $E$  is a semistable vector bundle over  $X$  (of any degree), then  $F_*E$  is also semistable.*

Unfortunately we do not know whether also stability is preserved by direct image under Frobenius. It has been shown that  $F_*L$  is stable for a line bundle  $L$  ([LanP] Proposition 1.2) and that in small characteristics the bundle  $F_*E$  is stable for any stable bundle  $E$  of small rank [JRXY]. The main ingredient of the proof is Faltings' cohomological criterion of semistability. We also need the fact that the generalized Verschiebung  $V$ , defined as the rational map from the moduli space  $\mathcal{M}_{X_1}(r)$  of semistable rank- $r$  vector bundles over  $X_1$  with fixed trivial determinant to the moduli space  $\mathcal{M}_X(r)$  induced by pull-back under the relative Frobenius map  $F$ ,

$$V_r : \mathcal{M}_{X_1}(r) \dashrightarrow \mathcal{M}_X(r), \quad E \longmapsto F^*E$$

is dominant for large  $r$ . We actually show a stronger statement for large  $r$ .

**1.2. Proposition.** *If  $l \geq g(p-1) + 1$  and  $l$  prime, then the generalized Verschiebung  $V_l$  is generically étale for any curve  $X$ . In particular  $V_l$  is separable and dominant.*

As an application of Theorem 1.1 we obtain an upper bound of the rational invariant  $\nu$  of a vector bundle  $E$ , defined as

$$\nu(E) := \mu_{\max}(F^*E) - \mu_{\min}(F^*E),$$

where  $\mu_{\max}$  (resp.  $\mu_{\min}$ ) denotes the slope of the first (resp. last) piece in the Harder-Narasimhan filtration of  $F^*E$ .

**1.3. Proposition.** *For any semistable rank- $r$  vector bundle  $E$*

$$\nu(E) \leq \min((r-1)(2g-2), (p-1)(2g-2)).$$

---

2000 *Mathematics Subject Classification.* Primary 14H40, 14D20, Secondary 14H40.

We note that the inequality  $\nu(E) \leq (r-1)(2g-2)$  was proved in [SB] Corollary 2 and in [S] Theorem 3.1. We suspect that the relationship between both inequalities comes from the conjectural fact that the length (=number of pieces) of the Harder-Narasimhan filtration of  $F^*E$  is at most  $p$  for semistable  $E$ .

Finally we show that direct images of line bundles under Frobenius are characterized by maximality of the invariant  $\nu$ .

**1.4. Proposition.** *Let  $E$  be a stable rank- $p$  vector bundle over  $X$ . Then the following statements are equivalent.*

- (1) *There exists a line bundle  $L$  such that  $E = F_*L$ .*
- (2)  $\nu(E) = (p-1)(2g-2)$ .

We do not know whether the analogue of this proposition remains true for higher rank.

## 2. REDUCTION TO THE CASE $\mu(E) = g - 1$ .

In this section we show that it is enough to prove Theorem 1.1 for semistable vector bundles  $E$  with slope  $\mu(E) = g - 1$ .

Let  $E$  be a semistable vector bundle over  $X$  of rank  $r$  and let  $s$  be the integer defined by the equality

$$\mu(E) = g - 1 + \frac{s}{r}.$$

Applying the Grothendieck-Riemann-Roch theorem to the Frobenius map  $F : X \rightarrow X_1$ , we obtain

$$\mu(F_*E) = g - 1 + \frac{s}{pr}.$$

Let  $\pi : \tilde{X} \rightarrow X$  be a connected étale covering of degree  $n$  and let  $\pi_1 : \tilde{X}_1 \rightarrow X_1$  denote its twist by the Frobenius of  $k$  (see [R] section 4). The diagram

$$(2.1) \quad \begin{array}{ccc} \tilde{X} & \xrightarrow{F} & \tilde{X}_1 \\ \pi \downarrow & & \downarrow \pi_1 \\ X & \xrightarrow{F} & X_1 \end{array}$$

is Cartesian and we have an isomorphism

$$\pi_1^*(F_*E) \cong F_*(\pi^*E).$$

Since semistability is preserved under pull-back by a separable morphism of curves, we see that  $\pi^*E$  is semistable. Moreover if  $F_*(\pi^*E)$  is semistable, then  $F_*E$  is also semistable.

Let  $L$  be a degree  $d$  line bundle over  $\tilde{X}_1$ . The projection formula

$$F_*(\pi^*E \otimes F^*L) = F_*(\pi^*E) \otimes L$$

shows that semistability of  $F_*(\pi^*E)$  is equivalent to semistability of  $F_*(\pi^*E \otimes F^*L)$ .

Let  $\tilde{g}$  denote the genus of  $\tilde{X}$ . By the Riemann-Hurwitz formula  $\tilde{g} - 1 = n(g - 1)$ . We compute

$$\mu(\pi^*E \otimes F^*L) = n(g - 1) + n\frac{s}{r} + pd = \tilde{g} - 1 + n\frac{s}{r} + pd,$$

which gives

$$\mu(F_*(\pi^*E \otimes F^*L)) = \tilde{g} - 1 + n\frac{s}{pr} + d.$$

**2.1. Lemma.** *For any integer  $m$  there exists a connected étale covering  $\pi : \tilde{X} \rightarrow X$  of degree  $n = p^k m$  for some  $k$ .*

*Proof.* If the  $p$ -rank of  $X$  is nonzero, the statement is clear. If the  $p$ -rank is zero, we know by Corollaire 4.3.4 [R] that there exist connected étale coverings  $Y \rightarrow X$  of degree  $p^t$  for infinitely many integers  $t$  (more precisely for all  $t$  of the form  $(l-1)(g-1)$  where  $l$  is a large prime). Now we decompose  $m = p^s u$  with  $p$  not dividing  $u$ . We then take a covering  $Y \rightarrow X$  of degree  $p^t$  with  $t \geq s$  and a covering  $\tilde{X} \rightarrow Y$  of degree  $u$ .  $\square$

Now the lemma applied to the integer  $m = pr$  shows existence of a connected étale covering  $\pi : \tilde{X} \rightarrow X$  of degree  $n = p^k m$ . Hence  $n \frac{s}{pr}$  is an integer and we can take  $d$  such that  $n \frac{s}{pr} + d = 0$ .

To summarize, we have shown that for any semistable  $E$  over  $X$  there exists a covering  $\pi : \tilde{X} \rightarrow X$  and a line bundle  $L$  over  $\tilde{X}_1$  such that the vector bundle  $\tilde{E} := \pi^* E \otimes F^* L$  is semistable with  $\mu(\tilde{E}) = \tilde{g} - 1$  and such that semistability of  $F_* \tilde{E}$  implies semistability of  $F_* E$ .

### 3. PROOF OF THEOREM 1.1

In order to prove semistability of  $F_* E$  we shall use the cohomological criterion of semistability due to Faltings [F].

**3.1. Proposition** ([L] Théorème 2.4). *Let  $E$  be a rank- $r$  vector bundle over  $X$  with  $\mu(E) = g - 1$  and  $l$  an integer  $> \frac{r^2}{4}(g-1)$ . Then  $E$  is semistable if and only if there exists a rank- $l$  vector bundle  $G$  with trivial determinant such that*

$$h^0(X, E \otimes G) = h^1(X, E \otimes G) = 0.$$

Moreover if the previous condition holds for one bundle  $G$ , it holds for a general bundle by upper semicontinuity of the function  $G \mapsto h^0(X, E \otimes G)$ .

**Remark.** The proof of this proposition (see [L] section 2.4) works over any algebraically closed field  $k$ .

By Proposition 1.2 (proved in section 4) we know that  $V_l$  is dominant when  $l$  is a large prime number. Hence a general vector bundle  $G \in \mathcal{M}_X(l)$  is of the form  $F^* G'$  for some  $G' \in \mathcal{M}_{X_1}(l)$ . Consider a semistable  $E$  with  $\mu(E) = g - 1$ . Then by Proposition 3.1  $h^0(X, E \otimes G) = 0$  for general  $G \in \mathcal{M}_X(l)$ . Assuming  $G$  general, we can write  $G = F^* G'$  and we obtain by adjunction

$$h^0(X, E \otimes F^* G') = h^0(X_1, F_* E \otimes G') = 0.$$

This shows that  $F_* E$  is semistable by Proposition 3.1.

### 4. PROOF OF PROPOSITION 1.2

According to [MS] section 2 it will be enough to prove the existence of a stable vector bundle  $E \in \mathcal{M}_{X_1}(l)$  satisfying  $F^* E$  stable and

$$h^0(X_1, B \otimes \text{End}_0(E)) = 0,$$

because the vector space  $H^0(X_1, B \otimes \text{End}_0(E))$  can be identified with the kernel of the differential of  $V_l$  at the point  $E \in \mathcal{M}_{X_1}(l)$ . Here  $B$  denotes the sheaf of locally exact differentials over  $X_1$  (see [R] section 4).

Let  $l \neq p$  be a prime number and let  $\alpha \in JX_1[l] \cong JX[l]$  be a nonzero  $l$ -torsion point. We denote by

$$\pi : \tilde{X} \rightarrow X \quad \text{and} \quad \pi_1 : \tilde{X}_1 \rightarrow X_1$$

the associated cyclic étale cover of  $X$  and  $X_1$  and by  $\sigma$  a generator of the Galois group  $\text{Gal}(\tilde{X}/X) = \mathbb{Z}/l\mathbb{Z}$ . We recall that the kernel of the Norm map

$$\text{Nm} : J\tilde{X} \longrightarrow JX$$

has  $l$  connected components and we denote by

$$i : P := \text{Prym}(\tilde{X}/X) \subset J\tilde{X}$$

the associated Prym variety, i.e., the connected component containing the origin. Then we have an isogeny

$$\pi^* \times i : JX \times P \longrightarrow J\tilde{X}$$

and taking direct image under  $\pi$  induces a morphism

$$P \longrightarrow \mathcal{M}_X(l), \quad L \longmapsto \pi_* L.$$

Similarly we define the Prym variety  $P_1 \subset JX_1$  and the morphism  $P_1 \rightarrow \mathcal{M}_{X_1}(l)$  (obtained by twisting with the Frobenius of  $k$ ). Note that  $\pi_{1*}L$  is semistable for any  $L \in P_1$  and stable for general  $L \in P_1$  (see e.g. [B]). Since  $F^*(\pi_{1*}L) \cong \pi_*(F^*L)$  — see diagram (2.1) — and since  $F^*$  induces the Verschiebung  $V_P : P_1 \rightarrow P$ , which is surjective, we obtain that  $\pi_{1*}L$  and  $F^*(\pi_{1*}L)$  are stable for general  $L \in P_1$ .

Therefore Proposition 1.2 will immediately follow from the next Proposition.

**4.1. Proposition.** *If  $l \geq g(p-1) + 1$  then there exists a cyclic degree  $l$  étale cover  $\pi_1 : \tilde{X}_1 \rightarrow X_1$  with the property that*

$$h^0(X_1, B \otimes \text{End}_0(\pi_{1*}L)) = 0$$

for general  $L \in P_1$ .

*Proof.* By relative duality for the étale map  $\pi_1$  we have  $(\pi_{1*}L)^* \cong \pi_{1*}L^{-1}$ . Therefore

$$\text{End}(\pi_{1*}L) \cong \pi_{1*}L \otimes \pi_{1*}L^{-1} \cong \pi_{1*}(L^{-1} \otimes \pi_1^* \pi_{1*}L)$$

by the projection formula. Moreover since  $\pi_1$  is Galois étale we have a direct sum decomposition

$$\pi_1^* \pi_{1*}L \cong \bigoplus_{i=0}^{l-1} (\sigma^i)^* L.$$

Putting these isomorphisms together we find that

$$\begin{aligned} H^0(X_1, B \otimes \text{End}(\pi_{1*}L)) &= H^0(X_1, B \otimes \pi_{1*}(\bigoplus_{i=0}^{l-1} L^{-1} \otimes (\sigma^i)^* L)) \\ &= \bigoplus_{i=0}^{l-1} H^0(X_1, B \otimes \pi_{1*}(L^{-1} \otimes (\sigma^i)^* L)) \\ &= H^0(X_1, B \otimes \pi_{1*} \mathcal{O}_{\tilde{X}_1}) \oplus \bigoplus_{i=1}^{l-1} H^0(X_1, B \otimes \pi_{1*}(L^{-1} \otimes (\sigma^i)^* L)). \end{aligned}$$

Moreover  $\pi_* \mathcal{O}_{\tilde{X}_1} = \bigoplus_{i=0}^{l-1} \alpha^i$ , which implies that

$$(4.1) \quad H^0(X_1, B \otimes \text{End}_0(\pi_{1*}L)) = \bigoplus_{i=1}^{l-1} H^0(X_1, B \otimes \alpha^i) \oplus \bigoplus_{i=1}^{l-1} H^0(X_1, B \otimes \pi_{1*}(L^{-1} \otimes (\sigma^i)^* L)).$$

Let us denote for  $i = 1, \dots, l-1$  by  $\phi_i$  the isogeny

$$\phi_i : P_1 \longrightarrow P_1, \quad L \longmapsto L^{-1} \otimes (\sigma^i)^* L.$$

Since the function  $L \mapsto h^0(X_1, B \otimes \text{End}_0(\pi_{1*}L))$  is upper semicontinuous, it will be enough to show the existence of a cover  $\pi_1 : \tilde{X}_1 \rightarrow X_1$  satisfying

- (1) for  $i = 1, \dots, l-1$ ,  $h^0(X_1, B \otimes \alpha^i) = 0$  (or equivalently,  $P$  is an ordinary abelian variety).
- (2) for  $M$  general in  $P$ ,  $h^0(X_1, B \otimes \pi_{1*}M) = 0$ .

Note that these two conditions imply that the vector space (4.1) equals  $\{0\}$  for general  $L \in P_1$ , because the  $\phi_i$ 's are surjective.

We recall that  $\ker(\pi_1^* : JX_1 \rightarrow J\tilde{X}_1) = \langle \alpha \rangle \cong \mathbb{Z}/l\mathbb{Z}$  and that

$$P_1[l] = P_1 \cap \pi_1^*(JX_1) \cong \alpha^\perp / \langle \alpha \rangle$$

where  $\alpha^\perp = \{\beta \in JX_1[l] \mid \omega(\alpha, \beta) = 1\}$  and  $\omega : JX_1[l] \times JX_1[l] \rightarrow \mu_l$  denotes the symplectic Weil form. Consider a  $\beta \in \alpha^\perp \setminus \langle \alpha \rangle$ . Then  $\pi_1^* \beta \in P_1[l]$  and

$$\pi_{1*} \pi_1^* \beta = \bigoplus_{i=0}^{l-1} \beta \otimes \alpha^i.$$

Again by upper semicontinuity of the function  $M \mapsto h^0(X_1, B \otimes \pi_{1*}M)$  one observes that the conditions (1) and (2) are satisfied because of the following lemma (take  $M = \pi_{1*}\beta$ ).

**4.2. Lemma.** *If  $l \geq g(p-1) + 1$  then there exists a pair  $(\alpha, \beta) \in JX_1[l] \times JX_1[l]$  satisfying*

- (1)  $\alpha \neq 0$  and  $\beta \in \alpha^\perp \setminus \langle \alpha \rangle$ ,
- (2) for  $i = 1, \dots, l-1$   $h^0(X_1, B \otimes \alpha^i) = 0$ ,
- (3) for  $i = 0, \dots, l-1$   $h^0(X_1, B \otimes \beta \otimes \alpha^i) = 0$ .

*Proof.* We adapt the proof of [R] Lemme 4.3.5. We denote by  $\mathbb{F}_l$  the finite field  $\mathbb{Z}/l\mathbb{Z}$ . Then there exists a symplectic isomorphism  $JX_1[l] \cong \mathbb{F}_l^g \times \mathbb{F}_l^g$ , where the latter space is endowed with the standard symplectic form. Note that composition is written multiplicatively in  $JX_1[l]$  and additively in  $\mathbb{F}_l^{2g}$ . A quick computation shows that the number of isotropic 2-planes in  $\mathbb{F}_l^g \times \mathbb{F}_l^g$  equals

$$N(l) = \frac{(l^{2g} - 1)(l^{2g-2} - 1)}{(l^2 - 1)(l - 1)}.$$

Let  $\Theta_B \subset JX_1$  denote the theta divisor associated to  $B$ . Then by [R] Lemma 4.3.5 the cardinality  $A(l)$  of the finite set  $\Sigma(l) := JX_1[l] \cap \Theta_B$  satisfies

$$A(l) \leq l^{2g-2}g(p-1).$$

Suppose that there exists an isotropic 2-plane  $\Pi \subset \mathbb{F}_l^g \times \mathbb{F}_l^g$  which contains  $\leq l-2$  points of  $\Sigma(l)$ . Then we can find a pair  $(\alpha, \beta)$  satisfying the 3 properties of the Lemma as follows: any nonzero point  $x \in \Pi$  determines a line ( $=\mathbb{F}_l$ -vector space of dimension 1). Since a line contains  $l-1$  nonzero points, we obtain at most  $(l-1)(l-2)$  nonzero points lying on lines generated by  $\Sigma(l) \cap \Pi$ . Since  $(l-1)(l-2) < l^2 - 1$  there exists a nonzero  $\alpha$  in the complement of these lines. Now we note that there are  $l-1$  affine lines parallel to the line generated by  $\alpha$  and the  $l$  points on any of these affine lines are of the form  $\beta\alpha^i$  for  $i = 0, \dots, l-1$  for some  $\beta \in \alpha^\perp \setminus \langle \alpha \rangle$ . The points  $\Sigma(l) \cap \Pi$  lie on at most  $l-2$  such affine lines, hence there exists at least one affine line parallel to  $\langle \alpha \rangle$  avoiding  $\Sigma(l)$ . This gives  $\beta$ .

Finally let us suppose that any isotropic 2-plane contains  $\geq l-1$  points of  $\Sigma(l)$ . Then we will arrive at a contradiction as follows: we introduce the set

$$S = \{(x, \Pi) \mid x \in \Pi \cap \Sigma(l) \text{ and } \Pi \text{ isotropic 2-plane}\}.$$

with cardinality  $|S|$ . Then by our assumption we have

$$(4.2) \quad |S| \geq (l-1)N(l).$$

On the other hand, since any nonzero  $x \in \mathbb{F}_l^g \times \mathbb{F}_l^g$  is contained in  $\frac{l^{2g-2}-1}{l-1}$  isotropic 2-planes, we obtain

$$(4.3) \quad |S| \leq \frac{l^{2g-2} - 1}{l - 1} A(l).$$

Putting (4.2) and (4.3) together, we obtain

$$A(l) \geq \frac{l^{2g} - 1}{l + 1}.$$

But this contradicts the inequality  $A(l) \leq l^{2g-2}g(p-1)$  if  $l \geq g(p-1) + 1$ . □

This completes the proof of Proposition 4.1. □

**Remark.** It has been shown [O] Theorem A.6 that  $V_r$  is dominant for any rank  $r$  and any curve  $X$ , by using a versal deformation of a direct sum of  $r$  line bundles.

**Remark.** We note that  $V_r$  is not separable when  $p$  divides the rank  $r$  and  $X$  is non-ordinary. In that case the Zariski tangent space at a stable bundle  $E \in \mathcal{M}_{X_1}(r)$  identifies with the quotient

$H^1(X_1, \text{End}_0(E))/\langle e \rangle$  where  $e$  denotes the nonzero extension class of  $\text{End}_0(E)$  by  $\mathcal{O}_{X_1}$  given by  $\text{End}(E)$ . Then the inclusion of homotheties  $\mathcal{O}_{X_1} \hookrightarrow \text{End}_0(E)$  induces an inclusion  $H^1(X_1, \mathcal{O}_{X_1}) \subset H^1(X_1, \text{End}_0(E))/\langle e \rangle$  and the restriction of the differential of  $V_r$  at the point  $E$  to  $H^1(X_1, \mathcal{O}_{X_1})$  coincides with the non-injective Hasse-Witt map.

### 5. PROOF OF PROPOSITION 1.3

Since we already know that  $\nu(E) \leq (r-1)(2g-2)$  ([SB], [S]) it suffices to show that  $\nu(E) \leq (p-1)(2g-2)$ .

We consider the quotient  $F^*E \rightarrow Q$  with minimal slope, i.e.,  $\mu(Q) = \mu_{\min}(F^*E)$  and  $Q$  semistable. By adjunction we obtain a nonzero morphism  $E \rightarrow F_*(Q)$ , from which we deduce (using Theorem 1.1) that

$$\mu(E) \leq \mu(F_*Q) = \frac{1}{p} (\mu_{\min}(F^*E) + (p-1)(g-1))$$

hence

$$\mu(F^*E) \leq \mu_{\min}(F^*E) + (p-1)(g-1).$$

Similarly we consider the subbundle  $S \hookrightarrow F^*E$  with maximal slope, i.e.,  $\mu(S) = \mu_{\max}(F^*E)$  and  $S$  semistable. Taking the dual and proceeding as above, we obtain that

$$\mu(F^*E) \geq \mu_{\max}(F^*E) - (p-1)(g-1).$$

Now we combine both inequalities and we are done.

**Remark.** We note that the inequality of Proposition 1.3 is sharp. The maximum  $(p-1)(2g-2)$  is obtained for the bundles  $E = F_*E'$  (see [JRXY] Theorem 5.3).

### 6. CHARACTERIZATION OF DIRECT IMAGES

Consider a line bundle  $L$  over  $X$ . Then the direct image  $F_*L$  is stable ([LanP] Proposition 1.2) and the Harder-Narasimhan filtration of  $F^*F_*L$  is of the form (see [JRXY])

$$0 = V_0 \subset V_1 \subset \dots \subset V_{p-1} \subset V_p = F^*F_*L, \quad \text{with} \quad V_i/V_{i-1} \cong L \otimes \omega_X^{p-i}.$$

In particular  $\nu(F_*L) = (p-1)(2g-2)$ . In this section we will show a converse statement.

More generally let  $E$  be a stable rank- $rp$  vector bundle with  $\mu(E) = g-1 + \frac{d}{rp}$  for some integer  $d$  and satisfying

- (1) the Harder-Narasimhan filtration of  $F^*E$  has  $l$  terms.
- (2)  $\nu(E) = (p-1)(2g-2)$ .

**Questions.** Do we have  $l \leq p$ ? Is  $E$  of the form  $E = F_*G$  for some rank- $r$  vector bundle  $G$ ? We will give a positive answer in the case  $r = 1$  (Proposition 6.1).

Let us denote the Harder-Narasimhan filtration by

$$0 = V_0 \subset V_1 \subset \dots \subset V_{l-1} \subset V_l = F^*E, \quad V_i/V_{i-1} = M_i.$$

satisfying the inequalities

$$\mu_{\max}(F^*E) = \mu(M_1) > \mu(M_2) > \dots > \mu(M_l) = \mu_{\min}(F^*E).$$

The quotient  $F^*E \rightarrow M_l$  gives via adjunction a nonzero map  $E \rightarrow F_*M_l$ . Since  $F_*M_l$  is semistable, we obtain that  $\mu(E) \leq \mu(F_*M_l)$ . This implies that  $\mu(M_l) \geq g-1 + \frac{d}{r}$ . Similarly taking the dual of the inclusion  $M_1 \subset F^*E$  gives a map  $F^*(E^*) \rightarrow M_1^*$  and by adjunction  $E^* \rightarrow F_*(M_1^*)$ . Let us denote  $\mu(M_1^*) = g-1 + \delta$ , so that  $\mu(F_*(M_1^*)) = g-1 + \frac{\delta}{p}$ . Because of semistability of  $F_*(M_1^*)$ , we obtain  $-(g-1 + \frac{d}{rp}) = \mu(E^*) \leq \mu(F_*(M_1^*))$ , hence  $\delta \geq -2p(g-1) - \frac{d}{r}$ . This implies that

$\mu(M_1) \leq (2p-1)(g-1) + \frac{d}{r}$ . Combining this inequality with  $\mu(M_l) \geq g-1 + \frac{d}{r}$  and the assumption  $\mu(M_1) - \mu(M_l) = (p-1)(2g-2)$ , we obtain that

$$\mu(M_1) = (2p-1)(g-1) + \frac{d}{r}, \quad \mu(M_l) = g-1 + \frac{d}{r}.$$

Let us denote by  $r_i$  the rank of the semistable bundle  $M_i$ . We have the equality

$$(6.1) \quad \sum_{i=1}^l r_i = rp.$$

Since  $E$  is stable and  $F_*(M_l)$  is semistable and since these bundles have the same slope, we deduce that  $r_l \geq r$ . Similarly we obtain that  $r_1 \geq r$ .

Note that it is enough to show that  $r_l = r$ . Since  $E$  is stable and  $F_*M_l$  semistable and since the two bundles have the same slope and rank, they will be isomorphic.

We introduce the integers for  $i = 1, \dots, l-1$

$$\delta_i = \mu(M_{i+1}) - \mu(M_i) + 2(g-1) = \mu(M_{i+1} \otimes \omega) - \mu(M_i).$$

Then we have the equality

$$(6.2) \quad \sum_{i=1}^{l-1} \delta_i = \mu(M_l) - \mu(M_1) + 2(l-1)(g-1) = 2(l-p)(g-1).$$

We note that if  $\delta_i < 0$ , then  $\text{Hom}(M_i, M_{i+1} \otimes \omega) = 0$ .

**6.1. Proposition.** *Let  $E$  be stable rank- $p$  vector bundle with  $\mu(E) = g-1 + \frac{d}{p}$  and  $\nu(E) = (p-1)(2g-2)$ . Then  $E = F_*L$  for some line bundle  $L$  of degree  $g-1+d$ .*

*Proof.* Let us first show that  $l = p$ . We suppose that  $l < p$ . Then  $\sum_{i=1}^{l-1} \delta_i = 2(l-p)(g-1) < 0$  so that there exists a  $k \leq l-1$  such that  $\delta_k < 0$ . We may choose  $k$  minimal, i.e.,  $\delta_i \geq 0$  for  $i < k$ . Then we have

$$(6.3) \quad \mu(M_k) > \mu(M_i) + 2(g-1) \quad \text{for } i > k.$$

We recall that  $\mu(M_i) \leq \mu(M_{k+1})$  for  $i > k$ . The Harder-Narasimhan filtration of  $V_k$  is given by the first  $k$  terms of the Harder-Narasimhan filtration of  $F^*E$ . Hence  $\mu_{\min}(V_k) = \mu(M_k)$ .

Consider now the canonical connection  $\nabla$  on  $F^*E$  and its first fundamental form

$$\phi_k : V_k \hookrightarrow F^*E \xrightarrow{\nabla} F^*E \otimes \omega_X \longrightarrow (F^*E/V_k) \otimes \omega_X.$$

Since  $\mu_{\min}(V_k) > \mu(M_i \otimes \omega)$  for  $i > k$  we obtain  $\phi_k = 0$ . Hence  $\nabla$  preserves  $V_k$  and since  $\nabla$  has zero  $p$ -curvature, there exists a subbundle  $E_k \subset E$  such that  $F^*E_k = V_k$ .

We now evaluate  $\mu(E_k)$ . By assumption  $\delta_i \geq 0$  for  $i < k$ . Hence

$$\mu(M_i) \geq \mu(M_1) - 2(i-1)(g-1) \quad \text{for } i \leq k,$$

which implies that

$$\deg(V_k) = \sum_{i=1}^k r_i \mu(M_i) \geq \text{rk}(V_k) \mu(M_1) - 2(g-1) \sum_{i=1}^k r_i (i-1).$$

Hence we obtain

$$p\mu(E_k) = \mu(V_k) \geq \mu(M_1) - 2(g-1)C,$$

where  $C$  denotes the fraction  $\frac{\sum_{i=1}^k r_i (i-1)}{\text{rk}(V_k)}$ . We will prove in a moment that  $C \leq \frac{p-1}{2}$ , so that we obtain by substitution

$$p\mu(E_k) \geq (2p-1)(g-1) + d - (g-1)(p-1) = p(g-1) + d = p\mu(E),$$

contradicting stability of  $E$ . Now let us show that  $C \leq \frac{p-1}{2}$  or equivalently

$$\sum_{i=1}^k ir_i \leq \frac{p+1}{2} \sum_{i=1}^k r_i.$$

But that is obvious if  $k \leq \frac{p-1}{2}$ . Now if  $k > \frac{p-1}{2}$  we note that passing from  $E$  to  $E^*$  reverses the order of the  $\delta_i$ 's, so that the index  $k^*$  for  $E^*$  satisfies  $k^* \leq \frac{p-1}{2}$ . This proves that  $l = p$ .

Because of (6.1) we obtain  $r_i = 1$  for all  $i$  and therefore  $E = F_*M_p$ . □

## 7. STABILITY OF $F_*E$ ?

Is stability also preserved by  $F_*$ ?

We show the following result in that direction.

**7.1. Proposition.** *Let  $E$  be a stable vector bundle over  $X$ . Then  $F_*E$  is simple.*

*Proof.* Using relative duality  $(F_*E)^* \cong F_*(E^* \otimes \omega_X^{1-p})$  we obtain

$$H^0(X_1, \text{End}(F_*E)) = H^0(X, F^*F_*E \otimes E^* \otimes \omega_X^{1-p}).$$

Moreover the Harder-Narasimhan filtration of  $F^*F_*E$  is of the form (see [JRXY])

$$0 = V_0 \subset V_1 \subset \dots \subset V_{p-1} \subset V_p = F^*F_*E, \quad \text{with} \quad V_i/V_{i-1} \cong E \otimes \omega_X^{p-i}.$$

We deduce that

$$H^0(X, F^*F_*E \otimes E^* \otimes \omega_X^{1-p}) = H^0(X, V_1 \otimes E^* \otimes \omega_X^{1-p}) = H^0(X, \text{End}(E)),$$

and we are done. □

## REFERENCES

- [B] A. Beauville: On the stability of the direct image of a generic vector bundle, preprint available at <http://math.unice.fr/~beauvill/pubs/imdir.pdf>
- [F] G. Faltings: Projective connections and G-bundles, *J. Alg. Geometry* 2, No. 3 (1993), 507-568
- [JRXY] K. Joshi, S. Ramanan, E.Z. Xia, J.K. Yu: On vector bundles destabilized by Frobenius pull-back, *Compositio Math.* 142 (2006), 616-630
- [L] J. Le Potier: Module des fibrés semi-stables et fonctions thêta, *Moduli of vector bundles* (Sanda 1994, Kyoto 1994) 83-101, *Lecture Notes in Pure and Appl. Math.* 179, Dekker, New York, 1996
- [LanP] H. Lange, C. Pauly: On Frobenius-destabilized rank-2 vector bundles over curves, [math.AG/0309456](http://math.AG/0309456)
- [LasP] Y. Laszlo, C. Pauly: The Frobenius map, rank 2 vector bundles and Kummer's quartic surface in characteristic 2 and 3, *Adv. Math.* 185, No. 2 (2004), 246-269
- [MS] V.B. Mehta, S. Subramanian: Nef line bundles which are not ample, *Math. Zeit.* 219 (1995), 235-244
- [O] B. Osserman: The generalized Verschiebung map for curves of genus 2, *Math. Ann.*, to appear
- [R] M. Raynaud: Sections des fibrés vectoriels sur une courbe, *Bull. Soc. Math. France*, Vol. 110 (1982), 103-125
- [SB] N.I. Shepherd-Barron: Semistability and reduction mod  $p$ , *Topology*, Vol. 37, No. 3 (1998), 659-664
- [S] X. Sun: Remarks on semistability of  $G$ -bundles in positive characteristic, *Compositio Math.* 119 (1999), No. 1, 41-52

TATA INSTITUTE OF FUNDAMENTAL RESEARCH, HOMI BHABHA ROAD, MUMBAI 400 005, INDIA  
*E-mail address:* vikram@math.tifr.res.in

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE MONTPELLIER II - CASE COURRIER 051, PLACE  
 EUGÈNE BATAILLON, 34095 MONTPELLIER CEDEX 5, FRANCE  
*E-mail address:* pauly@math.univ-montp2.fr