

A SMOOTH COUNTEREXAMPLE TO NORI'S CONJECTURE ON THE FUNDAMENTAL GROUP SCHEME

CHRISTIAN PAULY

ABSTRACT. We show that Nori's fundamental group scheme $\pi(X, x)$ does not base change correctly under extension of the base field for certain smooth projective ordinary curves X of genus 2 defined over a field of characteristic 2.

1. INTRODUCTION

In the paper [N] Madhav Nori introduced the fundamental group scheme $\pi(X, x)$ for a reduced and connected scheme X defined over an algebraically closed field k as the Tannaka dual group of the Tannakian category of essentially finite vector bundles over X . In characteristic zero $\pi(X, x)$ coincides with the étale fundamental group, but in positive characteristic it does not (see e.g. [MS]). By analogy with the étale fundamental group, Nori conjectured that $\pi(X, x)$ base changes correctly under extension of the base field. More precisely:

Nori's conjecture (see [MS] page 144 or [N] page 89) If K is an algebraically closed extension of k , then the canonical homomorphism

$$(1.1) \quad h_{X,K} : \pi(X_K, x) \longrightarrow \pi(X, x) \times_k \text{Spec}(K)$$

is an isomorphism.

In [MS] V.B. Mehta and S. Subramanian show that Nori's conjecture is false for a projective curve with a cuspidal singularity. In this note (Corollary 4.2) we show that certain *smooth* projective ordinary curves of genus 2 defined over a field of characteristic 2 also provide counterexamples to Nori's conjecture.

The proof has two ingredients: the first is an equivalent statement of Nori's conjecture in terms of F -trivial bundles due to V.B. Mehta and S. Subramanian (see section 2) and the second is the description of the action of the Frobenius map on rank-2 vector bundles over a smooth ordinary curve X of genus 2 defined over a field of characteristic 2 (see section 3). In section 4 we explicitly determine the set of F -trivial bundles over X .

I would like to thank V.B. Mehta for introducing me to these questions and for helpful discussions.

2. NORI'S CONJECTURE AND F -TRIVIAL BUNDLES

Let X be a smooth projective curve defined over an algebraically closed field k of characteristic $p > 0$. Let $F : X \rightarrow X$ denote the absolute Frobenius of X and F^n its n -th iterate for some positive integer n .

2.1. Definition. A rank- r vector bundle E over X is said to be F^n -trivial if

$$E \text{ stable} \quad \text{and} \quad F^{n*}E \cong \mathcal{O}_X^r.$$

2000 *Mathematics Subject Classification.* Primary 14H40, 14D20, Secondary 14H40.

2.2. Proposition ([MS] Proposition 3.1). *If the canonical morphism $h_{X,K}$ (1.1) is an isomorphism, then any F^n -trivial vector bundle E_K over $X_K := X \times_k \text{Spec}(K)$ is isomorphic to $E_k \otimes_k K$ for some F^n -trivial vector bundle E_k over X .*

3. THE ACTION OF THE FROBENIUS MAP ON RANK-2 VECTOR BUNDLES

We briefly recall some results from [LP1] and [LP2].

Let X be a smooth projective ordinary curve of genus 2 defined over an algebraically closed field k of characteristic 2. By [LP2] section 2.3 the curve X equipped with a level-2 structure can be uniquely represented by an affine equation of the form

$$(3.1) \quad y^2 + x(x+1)y = x(x+1)(ax^3 + (a+b)x^2 + cx + c),$$

for some scalars $a, b, c \in k$. Let \mathcal{M}_X denote the moduli space of S -equivalence classes of semistable rank-2 vector bundles with trivial determinant over X — see e.g. [LeP]. We identify \mathcal{M}_X with the projective space \mathbb{P}^3 (see [LP1] Proposition 5.1). We denote by $V : \mathbb{P}^3 \dashrightarrow \mathbb{P}^3$ the rational map induced by pull-back under the absolute Frobenius $F : X \rightarrow X$. There are homogeneous coordinates $(x_{00} : x_{01} : x_{10} : x_{11})$ on \mathbb{P}^3 such that the equations of V are given as follows (see [LP2] section 5)

$$(3.2) \quad V(x_{00} : x_{01} : x_{10} : x_{11}) = (\sqrt{abc}P_{00}^2(x) : \sqrt{b}P_{01}^2(x) : \sqrt{c}P_{10}^2(x) : \sqrt{a}P_{11}^2(x)),$$

with

$$\begin{aligned} P_{00}(x) &= x_{00}^2 + x_{01}^2 + x_{10}^2 + x_{11}^2, & P_{10}(x) &= x_{00}x_{10} + x_{01}x_{11}, \\ P_{01}(x) &= x_{00}x_{01} + x_{10}x_{11}, & P_{11}(x) &= x_{00}x_{11} + x_{10}x_{01}. \end{aligned}$$

Given a semistable rank-2 vector bundle E with trivial determinant, we denote by $[E] \in \mathcal{M}_X = \mathbb{P}^3$ its S -equivalence class. The semistable boundary of \mathcal{M}_X equals the Kummer surface Kum_X of X . Given a degree 0 line bundle N on X , we also denote the point $[N \oplus N^{-1}] \in \mathbb{P}^3$ by N .

3.1. Proposition ([LP1] Proposition 6.1 (4)). *The preimage $V^{-1}(N)$ of the point $N \in \text{Kum}_X \subset \mathcal{M}_X = \mathbb{P}^3$ with coordinates $(x_{00} : x_{01} : x_{10} : x_{11})$*

- *is a projective line, if $x_{00} = 0$.*
- *consists of the 4 square-roots of N , if $x_{00} \neq 0$.*

4. COMPUTATIONS

In this section we prove the following

4.1. Proposition. *Let $X = X_{a,b,c}$ be the smooth projective ordinary curve of genus 2 given by the affine model (3.1). Suppose that*

$$(4.1) \quad a^2 + b^2 + c^2 + a + c = 0.$$

Then there exists a nontrivial family $\mathcal{E} \rightarrow X \times S$ parametrized by a 1-dimensional variety S (defined over k) of F^4 -trivial rank-2 vector bundles with trivial determinant over X . Moreover any F^4 -trivial rank-2 vector bundle E with trivial determinant appears in the family \mathcal{E} , i.e., is of the form $(\text{id}_X \times s)^\mathcal{E}$ for some k -valued point $s : \text{Spec}(k) \rightarrow S$.*

We therefore obtain a counterexample to Nori's conjecture.

4.2. Corollary. *Let $X = X_{a,b,c}$ be a curve satisfying (4.1). Then for any algebraically closed extension K , the morphism $h_{X,K}$ is not an isomorphism*

Proof. Since S is 1-dimensional, there exists a K -valued point $s : \text{Spec}(K) \rightarrow S$, which is not a k -valued point. Then the bundle $E_K = (\text{id}_X \times s)^*\mathcal{E}$ over X_K is not of the form $E_k \otimes_k K$. Now apply Proposition 2.2. \square

Proof of Proposition 4.1. The method of the proof is to determine explicitly all F^n -trivial rank-2 vector bundles E over X for $n = 1, 2, 3, 4$. Taking tensor product of E with 2^{n+1} -torsion line bundles allows us to restrict attention to F^n -trivial vector bundles with trivial determinant.

We first compute the preimage under iterates of V of the point $A_0 \in \mathbb{P}^3$ determined by the trivial rank-2 vector bundle over X . We recall (see e.g. [LP1] Lemma 2.11 (i)) that the coordinates of $A_0 \in \mathbb{P}^3$ in the coordinate system $(x_{00} : x_{01} : x_{10} : x_{11})$ are $(1 : 0 : 0 : 0)$. It follows from Proposition 3.1 and equations (3.2) that $V^{-1}(A_0)$ consists of the 4 points

$$(4.2) \quad (1 : 0 : 0 : 0), \quad (0 : 1 : 0 : 0), \quad (0 : 0 : 1 : 0) \quad \text{and} \quad (0 : 0 : 0 : 1),$$

which correspond to the 2-torsion points of the Jacobian of X . Abusing notation we denote by A_1 both the 2-torsion line bundle on X and the point $(0 : 1 : 0 : 0) \in \mathbb{P}^3$.

Both points A_0 and A_1 correspond to S -equivalence classes of semistable rank-2 vector bundles. The set of isomorphism classes represented by the two S -equivalence classes A_0 and A_1 equal $\mathbb{P}\text{Ext}^1(A_1, A_1) \cup \{0\}$ and $\mathbb{P}\text{Ext}^1(\mathcal{O}_X, \mathcal{O}_X) \cup \{0\}$ respectively, where 0 denotes the trivial extensions $A_1 \oplus A_1$ and $\mathcal{O}_X \oplus \mathcal{O}_X$. Note that the two cohomology spaces $\text{Ext}^1(A_1, A_1)$ and $\text{Ext}^1(\mathcal{O}_X, \mathcal{O}_X)$ are canonically isomorphic to $H^1(\mathcal{O}_X)$. The pull-back by the absolute Frobenius F of X induces a rational map

$$F^* : \mathbb{P}\text{Ext}^1(A_1, A_1) \longrightarrow \mathbb{P}\text{Ext}^1(\mathcal{O}_X, \mathcal{O}_X),$$

which coincides with the projectivized p -linear map on the cohomology $H^1(\mathcal{O}_X) \rightarrow H^1(\mathcal{O}_X)$ induced by the Frobenius map F . Since we have assumed X ordinary, this p -linear map is bijective. Hence we obtain that there is only one (strictly) semistable bundle E such that $[E] = A_1$ and $F^*E \cong \mathcal{O}_X^2$, namely $E = A_1 \oplus A_1$. In particular there are no F^1 -trivial rank-2 vector bundles over X .

By Proposition 3.1 and using the equations (3.2), we easily obtain that the preimage $V^{-1}(A_1)$ is a projective line $\mathbb{L} \cong \mathbb{P}^1$, which passes through the two points

$$(1 : 1 : 1 : 1) \quad \text{and} \quad (0 : 0 : 1 : 1).$$

We now determine the bundles E satisfying $F^*E \cong A_1 \oplus A_1$. Given E with $[F^*E] = A_1 \in \mathbb{P}^3$ we easily establish the equivalence

$$F^*E \cong A_1 \oplus A_1 \quad \iff \quad \dim \text{Hom}(F^*E, A_1) = \dim \text{Hom}(E, F_*A_1) = 2.$$

Suppose that E is stable and $F^*E \cong A_1 \oplus A_1$. The quadratic map

$$\det : \text{Hom}(E, F_*A_1) \longrightarrow \text{Hom}(\det E, \det F_*A_1) = H^0(\mathcal{O}_X(w))$$

has nontrivial fibre over 0, since $\dim \text{Hom}(E, F_*A_1) = 2$. Hence there exists a nonzero $f \in \text{Hom}(E, F_*A_1)$ not of maximal rank. We consider the line bundle $N = \text{im } f \subset F_*A_1$. Since F_*A_1 is stable (see [LaP] Proposition 1.2), we obtain the inequalities

$$0 = \mu(E) < \deg N < \frac{1}{2} = \mu(F_*A_1),$$

a contradiction. Therefore E is strictly semistable and $[E] = [A_2 \oplus A_2^{-1}]$ for some 4-torsion line bundle A_2 with $A_2^{\otimes 2} = A_1$. The S -equivalence class $[A_2 \oplus A_2^{-1}]$ contains three isomorphism classes and a standard computation shows that only the decomposable bundle $A_2 \oplus A_2^{-1}$ is mapped by F^* to $A_1 \oplus A_1$. In particular there are no F^2 -trivial rank-2 bundles.

We now determine the coordinates of A_2 by intersecting the line \mathbb{L} , which can be parametrized by $(r : r : s : s)$ with $r, s \in k$, with the Kummer surface, whose equation is (see [LP2] Proposition 3.1)

$$c(x_{00}^2x_{10}^2 + x_{01}^2x_{11}^2) + b(x_{00}^2x_{01}^2 + x_{10}^2x_{11}^2) + a(x_{00}^2x_{11}^2 + x_{10}^2x_{01}^2) + x_{00}x_{01}x_{10}x_{11} = 0.$$

The computations are straightforward and will be omitted. Let $u \in k$ be a root of the equation

$$(4.3) \quad u^2 + u = b.$$

Then $u + 1$ is the other root. The coordinates of the two 4-torsion line bundles (modulo the canonical involution of the Jacobian of X) A_2 such that $A_2^{\otimes 2} = A_1$ are

$$(u : u : \sqrt{b} : \sqrt{b}) \quad \text{and} \quad (u + 1 : u + 1 : \sqrt{b} : \sqrt{b}).$$

Now the equation $u = 0$ (resp. $u + 1 = 0$) implies by (4.3) $b = 0$, which is excluded because we have assumed X smooth. So by Proposition 3.1 the preimage $V^{-1}(A_2)$ consists of the 4 line bundles A_3 such that $A_3^{\otimes 2} = A_2$. In particular there are no F^3 -trivial rank-2 bundles.

One easily verifies that the image under the rational map V given by (3.2) of the hyperplane $x_{00} = 0$ is the quartic surface given by the equation

$$(4.4) \quad bx_{11}^2x_{10}^2 + cx_{11}^2x_{01}^2 + ax_{10}^2x_{01}^2 + x_{00}x_{10}x_{01}x_{11} = 0.$$

When we replace $(x_{00} : x_{01} : x_{10} : x_{11})$ with $(u : u : \sqrt{b} : \sqrt{b})$ in (4.4) we obtain the equation

$$(4.5) \quad b^2 + u^2(1 + a + c) = 0.$$

Similarly replacing $(x_{00} : x_{01} : x_{10} : x_{11})$ with $(u + 1 : u + 1 : \sqrt{b} : \sqrt{b})$ in (4.4) we obtain the equation

$$(4.6) \quad b^2 + (u^2 + 1)(1 + a + c) = 0.$$

Finally the product of (4.5) with (4.6) equals (here one uses (4.3)) equation (4.1) up to a factor b^2 , which we can drop since $b \neq 0$ — note that we have assumed X smooth, hence $b \neq 0$ by [LP2] Lemma 2.1. To summarize we have shown that if (4.1) holds, then by Proposition 3.1 there exists an 8-torsion line bundle A_3 with $A_3^{\otimes 4} = A_1$ and such that the preimage $V^{-1}(A_3)$ is a projective line $\Delta \subset \mathbb{P}^3$.

Consider a point $[E] \in \Delta$ away from the Kummer surface — note that Δ is not contained in the Kummer surface Kum_X because its intersection is contained in the set of 16-torsion points. Then E is stable and $[F^*E] = [A_3 \oplus A_3^{-1}]$. There are three isomorphism classes represented by the S -equivalence class $[A_3 \oplus A_3^{-1}]$, namely the trivial extension $A_3 \oplus A_3^{-1}$ and two nontrivial extensions (for the details see [LP1] Remark 6.2). Since E is invariant under the hyperelliptic involution we obtain $F^*E = A_3 \oplus A_3^{-1}$ and finally that E is F^4 -trivial. Hence any stable point on Δ is F^4 -trivial.

Therefore, assuming (4.1), there exists a 1-dimensional subvariety $\Delta_0 \subset \mathcal{M}_X \setminus \text{Kum}_X$ parametrizing all F^4 -trivial rank-2 bundles. Passing to an étale cover $S \rightarrow \Delta_0$ ensures existence of a “universal” family $\mathcal{E} \rightarrow X \times S$ and we are done. □

Remark. Note that equation (4.1) depends on the choice of a nontrivial 2-torsion line bundle A_1 . If one chooses the 2-torsion line bundle $(0 : 0 : 1 : 0)$ or $(0 : 0 : 0 : 1)$ — see (4.2) — the corresponding equations are

$$a^2 + b^2 + c^2 + a + b = 0 \quad \text{or} \quad a^2 + b^2 + c^2 + b + c = 0.$$

REFERENCES

- [LaP] H. Lange, C. Pauly: On Frobenius-destabilized rank-2 vector bundles over curves, math.AG/0309456
- [LP1] Y. Laszlo, C. Pauly: The action of the Frobenius map on rank 2 vector bundles in characteristic 2, J. Alg. Geometry 11 (2002), 219-243
- [LP2] Y. Laszlo, C. Pauly: The Frobenius map, rank 2 vector bundles and Kummer’s quartic surface in characteristic 2 and 3, Adv. Math. 185, No. 2 (2004), 246-269
- [LeP] J. Le Potier: Lectures on vector bundles, Cambridge Studies in Advanced Mathematics 54, Cambridge University Press, 1997
- [MS] V.B. Mehta, S. Subramanian: On the fundamental group scheme, Invent. Math. 148 (2002), 143-150

[N] M.V. Nori, The fundamental group scheme, Proc. Indian Acad. Sci. (Math. Sci.) 91 (1982), 78-122

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ DE MONTPELLIER II - CASE COURRIER 051, PLACE
EUGÈNE BATAILLON, 34095 MONTPELLIER CEDEX 5, FRANCE

E-mail address: `pauly@math.univ-montp2.fr`